

Grade 8

NUMBER SENSE AND NUMERATION: SQUARE ROOTS

This resource may be copied in its entirety, but is **not to be used for commercial purposes** without permission from the Centre for Education in Mathematics and Computing, University of Waterloo.

Play **Square Root Tic-Tac-Toe** first.

Click on <http://www.funbrain.com/cgi-bin/ttt.cgi?A1=s&A2=17&A3=0>

Read the **Lesson on Square Roots** at <http://argyll.epsb.ca/jreed/math8/strand1/1105.htm#17>

You can go to www.wiredmath.ca for the links.

The square root of 16 is written as $\sqrt{16}$. The value of $\sqrt{16}$ is 4 since $(4) \times (4) = 16$.

The **square root** of a number is one of its two *equal* factors.

For example, $\sqrt{144} = 12$ since $(12) \times (12) = 144$.

The symbol $\sqrt{\quad}$ is called a radical sign. The number under a radical sign is called the radicand. For example, for $\sqrt{5}$, 5 is the radicand.

1. Determine each square root.

a. $\sqrt{25}$

b. $\sqrt{49}$

c. $\sqrt{121}$

d. $\sqrt{9}$

e. $\sqrt{0}$

f. $\sqrt{64}$

g. $\sqrt{36}$

h. $\sqrt{4}$

i. $\sqrt{81}$

j. $\sqrt{100}$

k. $\sqrt{1}$

l. $\sqrt{256}$

m. $\sqrt{225}$

n. $\sqrt{169}$

o. $\sqrt{196}$

p. $\sqrt{400}$

q. $\sqrt{625}$

r. $\sqrt{1024}$

Note: $\sqrt{\frac{9}{4}} = \frac{\sqrt{9}}{\sqrt{4}} = \frac{3}{2}$

2. Determine each square root.

a. $\sqrt{\frac{49}{36}}$

b. $\sqrt{\frac{121}{64}}$

c. $\sqrt{\frac{25}{16}}$

d. $\sqrt{\frac{100}{16}}$

e. $\sqrt{\frac{144}{225}}$

f. $\frac{\sqrt{0}}{\sqrt{81}}$

g. $\sqrt{\frac{9 \times 9}{3 \times 3}}$

h. $\sqrt{\frac{12 \times 12}{6 \times 6}}$

i. $\sqrt{\frac{10 \times 10}{4 \times 4}}$

j. $\sqrt{\frac{8 \times 8}{12 \times 12}}$

k. $\sqrt{\frac{2 \times 2}{3 \times 3}}$

l. $\sqrt{\frac{6 \times 6}{7 \times 7}}$

Expectations: i) represent, compare, and order square roots; ii) estimate, and verify using a calculator, the positive square roots of whole numbers, and distinguish between whole numbers that have whole-number square roots (i.e., perfect square numbers) and those that do not.

For more activities and resources from the University of Waterloo's Faculty of Mathematics, please visit www.cemc.uwaterloo.ca.

The square root of a positive integer that is not a perfect square is an approximate value.

For example, $\sqrt{54}$ is close to $\sqrt{49}$ which equals 7.

Using a calculator $\sqrt{54} \doteq 7.3$, correct to one decimal place.

3. a. Estimate each square root without a calculator.
b. Use a calculator to determine each square root correct to one decimal place.
- i. $\sqrt{12}$ ii. $\sqrt{18}$ iii. $\sqrt{29}$ iv. $\sqrt{72}$ v. $\sqrt{83}$
vi. $\sqrt{119}$ vii. $\sqrt{0.25}$ viii. $\sqrt{1.44}$ ix. $\sqrt{2.25}$ x. $\sqrt{0.9}$

4. It can be shown that from a height of h metres, a person can see a distance d kilometres to the horizon, where $d = 3.53\sqrt{h}$. When the elevator of the CN Tower reaches the 200 m height, how far can the passengers in the elevator see across Lake Ontario?
5. Determine the length of a side of a square that has an area of 100 cm^2 .
6. Two sides of rectangles are given.
a. Determine the area of each rectangle.
b. Determine the length of a side of a square with the same area as the rectangle.

i. 2 cm and 8 cm

ii. 8 cm and 18 cm

iii. 8 cm and 25 cm

Did You Know?


The process of putting together the top antenna of the CN Tower took more than 3.5 weeks.



A Slice of History

Pythagoras (puh-thag-or-us) was a Greek philosopher and mathematician born in Samos. He and his followers tried to explain everything with numbers.

Pythagoras, 580 B.C. – 500 B.C. determined the relationship among the sides of a right-angled triangle. He concluded that the area of the square on the hypotenuse is equal to the sum of the areas of the squares on the other two sides.




Expectations: i) represent, compare, and order square roots; ii) estimate, and verify using a calculator, the positive square roots of whole numbers, and distinguish between whole numbers that have whole-number square roots (i.e., perfect square numbers) and those that do not.

For more activities and resources from the University of Waterloo's Faculty of Mathematics, please visit www.cemc.uwaterloo.ca.

The Pythagorean Theorem to find an unknown side of a right triangle

To find the length of the hypotenuse:




$$x^2 = 3^2 + 4^2$$

$$x^2 = 9 + 16$$

$$x = \sqrt{25}$$

$$x = 5$$

To find the length of a side when the hypotenuse is given:









$$5^2 + b^2 = 13^2$$

$$b^2 = 169 - 25$$







$$b = \sqrt{144}$$

$$b = 12$$

7. Determine the length of each unknown side of the following right-angled triangles.



8. Find the length of a diagonal of each rectangle.



Don't forget now! Go to www.wiredmath.ca for the link.

TRY THESE!

Square Root of a Perfect Square Number

<http://www.quia.com/jg/65631.html>



Square Root Flashcards

<http://www.aplusmath.com/Flashcards/sqrt.html>

Expectations: i) represent, compare, and order square roots; ii) estimate, and verify using a calculator, the positive square roots of whole numbers, and distinguish between whole numbers that have whole-number square roots (i.e., perfect square numbers) and those that do not.

For more activities and resources from the University of Waterloo's Faculty of Mathematics, please visit www.cemc.uwaterloo.ca.

CHALLENGE YOURSELF

9. Without using a calculator, determine each value.

Write your answer as a fraction in the form $\frac{a}{b}$, $b \neq 0$.

a. $\sqrt{\frac{50}{32}}$


b. $\sqrt{\frac{128}{450}}$

c. $\sqrt{\frac{48}{147}}$

d. $\frac{\sqrt{45}}{\sqrt{125}}$

e. $\frac{\sqrt{288}}{\sqrt{200}}$

10. The cube, as shown, has a total surface area of 1176 cm^2 . Determine the length of one of its edges.



A Slice of History

Sometimes called Hero, **Heron of Alexandria** was an important geometer and worker in mechanics.


11. If a , b and c represent the lengths of the sides of any triangle, the area is given by

Heron's Formula $A = \sqrt{s(s-a)(s-b)(s-c)}$ where $s = \frac{a+b+c}{2}$.


- a. Determine the exact area of a triangle with sides 7 cm, 24 cm and 25 cm. Discuss another method to find the area of this triangle. Why does the method work?
- b. Determine the area of a triangle with sides of length 16 cm, 16 cm and 8 cm. Round the answer off to the nearest unit.

EXTENSIONS

12. A square is inscribed in a circle, as shown. The area of the square is 98 cm^2 . Determine the radius of the circle.



13. Determine the length of x .



14. Observe that $1^3 + 2^3 = (1+2)^2$, $1^3 + 2^3 + 3^3 = (1+2+3)^2$ and $1^3 + 2^3 + 3^3 + 4^3 = (1+2+3+4)^2$. If the same pattern holds, then what is the value of $\sqrt{1^3 + 2^3 + 3^3 + 4^3 + 5^3 + 6^3 + 7^3 + 8^3 + 9^3 + 10^3}$?

Expectations: i) represent, compare, and order square roots; ii) estimate, and verify using a calculator, the positive square roots of whole numbers, and distinguish between whole numbers that have whole-number square roots (i.e., perfect square numbers) and those that do not.

For more activities and resources from the University of Waterloo's Faculty of Mathematics, please visit www.cemc.uwaterloo.ca.